# ICD-11 vs ICD-10-CM Coding Cheat Sheet

## DSM-5-TR → ICD-10-CM → ICD-11 crosswalk for common diagnoses

Side-by-side code structures for the diagnoses clinicians search most. ICD-11 uses stem codes + postcoordination extensions; the US still bills ICD-10-CM.

| Diagnosis | DSM-5-TR | ICD-10-CM | ICD-11 |
|---|---|---|---|
| Major Depressive Disorder (single) | F32.x | F32.x | 6A70 |
| Recurrent Depressive Disorder | F33.x | F33.x | 6A71 |
| Persistent Depressive Disorder | F34.1 | F34.1 | 6A72 |
| Bipolar I Disorder | F31.x | F31.x | 6A60 |
| Bipolar II Disorder | F31.81 | F31.81 | 6A61 |
| Cyclothymic Disorder | F34.0 | F34.0 | 6A62 |
| Generalized Anxiety Disorder | F41.1 | F41.1 | 6B00 |
| Panic Disorder | F41.0 | F41.0 | 6B01 |
| Agoraphobia | F40.0x | F40.0x | 6B02 |
| Specific Phobia | F40.2xx | F40.2xx | 6B03 |
| Social Anxiety Disorder | F40.1x | F40.1x | 6B04 |
| PTSD | F43.1x | F43.1x | 6B40 |
| Complex PTSD | — | F43.9 | 6B41 |
| Adjustment Disorder | F43.2x | F43.2x | 6B43 |
| Prolonged Grief Disorder | F43.8 | F43.8 | 6B42 |
| Obsessive-Compulsive Disorder | F42.x | F42.x | 6B20 |
| Hoarding Disorder | F42.3 | F42.3 | 6B24 |
| ADHD | F90.x | F90.x | 6A05 |
| Autism Spectrum Disorder | F84.0 | F84.0 | 6A02 |
| Specific Learning Disorder | F81.x | F81.x | 6A03 |
| Tourette Syndrome | F95.2 | F95.2 | 8A05.0 |

# Coding Cheat Sheet — continued

Remaining diagnoses + general medical crosswalk

| Diagnosis | DSM-5-TR | ICD-10-CM | ICD-11 |
|---|---|---|---|
| Schizophrenia | F20.x | F20.x | 6A20 |
| Schizoaffective Disorder | F25.x | F25.x | 6A21 |
| Delusional Disorder | F22 | F22 | 6A24 |
| Brief Psychotic Disorder | F23 | F23 | 6A23 |
| Anorexia Nervosa | F50.0x | F50.0x | 6B80 |
| Bulimia Nervosa | F50.2 | F50.2 | 6B81 |
| Binge Eating Disorder | F50.81 | F50.81 | 6B82 |
| Borderline Personality Disorder | F60.3 | F60.3 | 6D11 |
| Alcohol Use Disorder | F10.2x | F10.2x | 6C40 |
| Premenstrual Dysphoric Disorder | F32.81 | N94.3 | GA34.41 |
| Type 2 Diabetes Mellitus | — | E11.x | 5A11 |
| Essential Hypertension | — | I10 | BA00 |
| Asthma | — | J45.x | CA23 |
| Chronic Low Back Pain | — | M54.5 | MG30.02 |
| GERD | — | K21.9 | DA22 |
| Obesity | — | E66.9 | 5B81 |

## How to read the ICD-11 column

• ICD-11 codes are alphanumeric stems (e.g., 6A70) extended with /severity or /laterality suffixes — postcoordination replaces ICD-10's combined codes.

• Mental health moved to chapter 06 (6A00–6E8Z): mood 6A60–6A8Z, anxiety 6B00–6B0Z, OCD 6B20–6B2Z, stress 6B40–6B4Z, eating 6B80–6B8Z.

• Personality disorders are now dimensional (6D10–6D11) rather than categorical — borderline pattern codes to 6D11.

• US payers still require ICD-10-CM. CMS launched the ICD-11 Add-on Code Beta Nov 2024; no mandatory transition date yet.

• Verify against the current ICD-10-CM tabular list and CMS ICD-11 readiness notices before claim submission. This sheet is educational, not coding advice.